



ISSN: 2395-1958
IJOS 2019; 5(2): 899-902
© 2019 IJOS
www.orthopaper.com
Received: 19-02-2019
Accepted: 13-03-2019

Dr. Bhavesh R Namsha
Associate Professor, Dept. of
Orthopaedics, SSG hospital and
Medical College, Baroda,
Gujarat, India

Dr. Uday Meghnathi
Senior Resident, Dept. of
Orthopaedics, SSG hospital and
Medical College, Baroda,
Gujarat, India

Dr. Bhavik K Ahir
Junior Resident, Dept. of
Orthopaedics, SSG hospital and
Medical College, Baroda,
Gujarat, India

Dr. Alizayagam N Hasan
Senior Resident, Dept. of
Orthopaedics, SSG hospital and
Medical College, Baroda,
Gujarat, India

Correspondence

Dr. Uday Meghnathi
Senior Resident, Dept. of
Orthopaedics, SSG hospital and
Medical College, Baroda,
Gujarat, India

International Journal of Orthopaedics Sciences

Volar plating in distal end radius fractures and its clinical and radiological outcomes

Dr. Bhavesh R Namsha, Dr. Uday Meghnathi, Dr. Bhavik K Ahir and Dr. Alizayagam N Hasan

DOI: <https://doi.org/10.22271/ortho.2019.v5.i2m.108>

Abstract

Introduction: Fractures of distal end of radius are one of the most common skeletal injuries encountered in orthopaedics. There are various modalities for treatment like closed reduction and cast application, external fixators, ORIF with plate etc.

Materials and Methods: We conducted a prospective study of 20 patients having AO OTA type A, B & C distal radius fractures treated with volar locking plate at our institute between 2018 to 2019. We include all close injuries in skeletally mature patients between ages 20-65 years. Outcome of the study was evaluated by using the Gartland and Werley score.

Results: 5 patients had AO OTA type A, 8 patients had type B and 7 patients had type C fractures. Mean palmar flexion was 78 degrees. Dorsi flexion was 80 degrees observed. Average supination and pronation were 85 & 78 degrees respectively. Radial & ulnar deviations average 12 and 24 degrees were observed. More than 90% of the patients had excellent to good outcome based on Gartland and Werley score.

Conclusions: Volar plating is a proven method of choice in majority of distal end radius fractures regardless of comminution and fragments separation.

Keywords: Distal radius fractures, internal fixation, volar plate, AO OTA fractures

Introduction

Fractures of distal end of radius are one of the most common skeletal injuries encountered in orthopaedics [1]. They occur most frequently in patients after the fourth decade of life [2]. Road accidents play a major role sustaining these trauma while older persons has house fall with trivial insults [3, 4] Pathoanatomy & biomechanics of radiocarpal & radioulnar joints have pivotal role in maintaining wrist functions and movements [5]. Problems associated with these fractures hamper strength of grip and early arthritis with carpal instability.

Congruity of articular surfaces can only achieved with ligamentotaxis and reconstruction of joint addressing each fragment along with early rehabilitation.5 To date, valid components of treatment strategy includes reduction cast, ORIF with dorsal and volar approach locking plates and external fixators [6]. Geriatric patients with less functional demands can be treated with non invasive cast reductions [1] External fixators opt adequate ligamentotaxis but has inherent complications of pins with suboptimal articular reduction [2]. Dorsal barton fractures with comminutions where buttress needed, dorsal approach low profile plates have excellent offset. But since no hardware is escaped without trouble, extensor tendons may damage [7].

So volar plating is the mainstay of technique to achieve anatomic reductions and ligamentous reconstructions in majority of articular fractures of lower end of radius with less complications with meticulous surgeon [7].

Materials and Methods

In our study, we included 20 patients having lower end of radius fracture coming to our institute as outdoor or emergency department treated with volar locking plates between years 2018 to 2019. Among them 15 are male and 5 are female with age ranges from 20 years to 65years.

We use AO/OTA classification for distal radius fracture on the basis of plain radiographs. Patient's outcome measurements were determined using the Gartland & Werley score.

Inclusion criteria

1. AO-OTA type A, B, C with subgroups
2. Close distal end radius fractures
3. Skeletally mature patients.

Exclusion criteria

1. Fractures with diaphyseal involvement/isolated radial styloid fractures.
2. Associated fracture of carpal bones.
3. Delayed presentation >2weeks. Pathological fractures.

Initial assessment of all the patients done to rule out any other associated injuries and splinting of affected limb was done. Clinical and radiographic evaluation of fracture was done by obtaining plain radiographs to classify the fractures. Patients operated after proper counselling and informed consent. Operation done under regional or general anaesthesia using tourniquet and proper aseptic precautions.

Surgical technique: Skin incision (Henry approach) taken at volar aspect of wrist between flexor carpi radialis tendon and radial artery. Skin subcutaneous tissue cut and haemostasis achieved. By incising the tendon sheath and retracting the flexor polices longus to the ulnar side pronator quadratus is exposed. By cutting pronator quadratus in L shape manner and retracting it, volar surface of distal radius is exposed.

Pronator quadratus was repaired using absorbable sutures. Subcutaneous and skin closure was done. Postoperatively radiographs were taken, the limb was kept elevated in below elbow plaster slab, active finger, forearm rotation and shoulder exercises were started at the earliest possible. The plaster slab was removed within 2 weeks, crepe bandage applied and active exercises of wrist, elbow and shoulder were started. Heavy lifting was not allowed until signs of fracture healing were radiographically confirmed.

Reduction done by manual traction and hold temporarily by k wires if necessary under IITV guide. Appropriate size plate selected and placed over volar surface of radius and temporarily fixed with cortical screw under IITV guidance. After confirming the desired reduction, remaining screws are fixed under IITV guide.

Evaluation of outcome

Radio logically: The follow up protocol was 6 weeks, 3 months, and 6 months. Clinical & radiographic assessments were performed at every visit. Radiological assessment included measurements of volar tilt, radial height & radial inclination.

Clinical & functional: Wrist range of motion was measured in flexion, extension, pronation, supination, ulnar deviation & radial deviation and was compared with uninjured limb. The overall function of the upper limb was assessed using the Gartl and & Werleyscore.

Table 1: Scoring system (Gartland & Werley (G & W) Score)

G&W Score	Number of patients	Percentage
Excellent 0-2	17	85%
Good 3-8	2	10%
Fair 9-20	1	5%
Poor >20	0	0%
Total	20	100%

There were 95% of patients having excellent to good palmar flexion range at wrist at final follow up. One patient has fair

outcome and she presented as residual dorsal tilt at final follow up. Most of the patients had palmar flexion range from 65–85 degrees. The average palmar flexion was 78degrees.

All the patients have excellent to good results in terms of range of dorsi flexion at wrist at final follow up. All the patients had dorsi flexion range from 60- 85 degrees. The average dorsi flexion was 80 degrees. There were 96% of patients having excellent to good supination range at wrist joint at final follow up. All the patients except one had supination range from 76-85 degrees. The average supination was 85degrees. All the patients have excellent to good pronation range at wrist joint at final follow up. The average pronation at final follow up was 78 degrees.

In our study, 95% of the patients had excellent to good radial deviation range at wrist at final follow up. Most of the patients (18) having radial deviation range from 10-20 degrees. The average radial deviation was 12 degrees.

All the patients in our study had excellent to good ulnar deviation range at wrist at final follow up. All of the patients having ulnar deviation range of 20-30 degrees. The average ulnar deviation at final follow up was 24 degrees.

X-rays of the patient were evaluated for Radiological assessment of Radial inclination, Radial shortening and Palmer tilt evaluated according to.

In our study, more than 80% of the patients had loss of radial inclination less than 9 degrees and had excellent to good results. The average radial inclination at final follow up was 19 degrees.

In this study of 20 patients, all the patients had less than 6 mm radial shortening and had excellent to good results at final follow up. The average radial length at final follow up was 10 mm. More than 60% patients had excellent to good results with regard to loss of palmar tilt less than 6 degrees. There were 9 patients showing loss of palmar tilt upto 14 degrees and had fair outcome. It could be due to dorsal comminution and dorsal collapse on subsequent follow up. Average palmar tilt at final follow up was around 7degrees.

More than 90% of the patients had excellent to good outcome based on Gartland and Werley score. Among them one patient had fair outcome. That patient had dorsal tilt at final follow up. We observed prominence at ulnar styloid in 2 patients (10%), residual dorsal tilt in 1 patients (5%), reflex sympathetic dystrophy in 1(5%) patient and hand grip strength weakness in 1(5%) patient. The distal locking screws have to be placed as close to the articular surface as possible. These screws act as a rake, maintaining reduction. Some oblique intraoperative views are recommended to precisely assess screws tips penetration and trajectory^[11].

We didn't observe tendon related injuries in our series of patients as flexor tendons are vulnerable to injury if plates are fixed beyond topographical boundary which in turn leads to flexor tendon synovitis. In our series no extensor tendons related problems encountered as these tendons are frequently irritated by longer tips of locking screws from plates fixed volarly. Some of the patients present with complex regional pain syndrome. This may be due to prolong immobilization and inadequate physiotherapy. Study conducted on large series of patients Kenny Kwan *et al.*^[10] showed^[11] complications at various stages of follow-ups postoperatively. We have come across some of the complications. Among all,^[2] patients having prominent ulnar styloid process and 1 patient having residual dorsal tilt. One patient developed RSD. One patient had loss of reduction subsequent follow up. None of our patient got infected.

Jupiter *et al*^[8] study advocated mean Gartland and Werley

score improved significantly from 4 points at six months to 2 points at two years.

We achieved Gartland and Werley score excellent results in 85% and good results in 10% of patients. Denju Osada *et al.* [14] and Kenny Kwan *et al.* [10] demonstrated 96% and 91% excellent results respectively.

Discussion

Chung *et al.* [9] hypothesized demographical and population based incidence of internal fixation of distal end radius is on disposal. We have included 20 patients, among them 15 were male and 5 were female with an average age of 39 years (20–65 years) having distal end of radius fracture treated with volar locking plate from 2018 to 2019 in our institute. We observed mean age in our study was 39 years. In a study by Kenny Kwan *et al.* [10] mean age was 51 years.

In a study conducted by Kenny Kwan *et al.* [10] mean supination and pronation were 86 and 80, we observed clinically 85 degrees supination and 78 degrees pronation. We have small sample size compared to this study which has sample size 75 patients. Radial and ulnar deviations are 12 and 24 degrees in our study while 23 and 37 degrees respectively in Denju Osada *et al.* [14] series.

In our study the radiological outcome are within acceptable criteria. Clinical outcome in view of wrist range of motion and grip strength are also satisfactory. The total functional result on the basis of Gartland and Werley score was 85% of excellent and 10% of good outcome on final follow up at approximate 6 month of follow up.

The primary goal in treatment of unstable fractures of the distal radius is to achieve proper reconstruction of the disrupted anatomy and expedite return of hand function without complications. Volar plating is considered superior in clinical radiological and surgical outcome of patients compared to dorsal plating in majority of distal end radius AO type fractures.

Palmar flexion in our study mean 78 degree achieved while in Denju Osada *et al.* series 66 degree achieved. Wrist dorsiflexion mean 75 degrees was in a study by Denju *et al.* while we achieved mean 80 degrees of dorsiflexion

Surgical approach, fracture reduction, and volar plating techniques are easier than dorsal plating. Vascularity of dorsal distal radius is not hampered with volar plating. Extensor tendon handling in dorsal approach may injure these structures which is preserved in palmar approach. Closure of the wound preceded by well coverage of healthy pronator quadratus breasting over volar plate.

With the advent of low profile fragment specific and other locking plates flexor tendons are very well preserved and safeguarded by volar plating if placed meticulously just before watershed line. In absence of severe comminutions, volar plating confers desirable successful articular reduction and satisfactory outcome.

Anatomical reduction of the palmar cortex may avoid the shortening of the radius, which is important to its restoration. Plate requires bending according to AO principle of fracture reduction despite of relatively flat volar surface compared to dorsal distal radius [13, 14, 11].

Sobky *et al.* [12] biomechanical study indicates volar fixation of unstable distal radius fractures with a fixed angle device is a reliable means of stabilization.

Most complications of distal radial fracture Treatment are the consequence of required immobilization of the wrist for between 4 and 6 weeks with either a plaster cast or an external fixator. This also applies to operative methods either by volar

plating with non-locking screws, combined volar and dorsal plating AO OTA type A and B fractures with less comminution rarely need a period of immobilization although type C also don't require immobilization fixed with latest locking low profile fragment specific plates [11].

In order to achieve reproducibly good results, some aspects of the surgical technique are very important.

Conclusion

Accurate anatomical reduction of distal end radius fracture with internal fixation with volar locking plate is promising and proven technique. Volar plate fixation is mainstay in management of distal end radius fractures with good clinical outcome and limited complications. Flexor as well as extensor tendons are safeguarded by meticulous volar approach and by learning plate fixation techniques.

Conflict of interest: Nil

References

1. Kapoor H, Agarwal A, Dhaon BK. Displaced intra-articular fractures of distal radius: a comparative evaluation of results following closed reduction, external fixation and open reduction with internal fixation. *Injury*. 2000; 31(2):75-9.
2. Caporrino FA, Belotti JC, Ulson HJR, Toledo LFQ, Reis FB, Machado JKS. Fraturas da extremidade distal do rádio e da ulna. In: Pardini Júnior AG, Freitas A. *Traumatisms da mão*. Rio de Janeiro: Med Book. 2008; 4:411-45.
3. Joideep Phadnis, Alex Trompeter, Kirren Gallagher, Lucy Bradshaw, David S Elliott, Kevin Newman. Mid-term functional outcome after the internal fixation of distal radius fractures. *Journal of Orthopedics Surgery and Research*. 2012; 7:4.
4. Mader K, Pennig D. The Treatment of Severely Comminuted intra articular fractures of distal radius, *Strat Trauma Limb Recon*. 2006; 1:2-17.
5. Wong KK, Chan KW, Kwok TK, Mak KH. Volar fixation of dorsally displaced distal radius fractures using locking compression plate, *Journal of Orthopedics Surgery*. 2005; 13(2):153-157.
6. Michele Rampoldi, Dante Palombi, Dontella Tagliente. Distal Radius Fractures with Diaphyseal Involvement: Fixation With fixed Angle Volar Plate. *J Orthopaed T Raumatol*. 2011; 12:137-143.
7. Jorge Orbay. Volar plate fixation of distal radius fractures. *Hand clinics*. 2005; 21:347-354.
8. Jesse B Jupiter, Marent Huber M. and LCP study group, Operation management of Distal Radial fractures with 2.4 mm locking plates. A multicenter prospective case series. *J Bone Joint Surg Am*. 2009; 91:55-65
9. Kevin C Chung, Melissa J Shauver, Huiying Yin, Myra Kim H, Onur Baser, John D Birkmeyer. Variation in the use of internal fixation of distal radius fracture in united states population. *J Bone Joint Surg Am*. 2011; 93:2154-2162.
10. Kenny Kwan, Tak Wing Lau, Frankie Leung. Operative treatment of distal radius fractures with locking plate system- A Prospective study. *International Orthopedics (SICOT)*. 2011; 35:389-394.
11. Drobetz H, Kutscha-Lissberg E. Osteosynthesis of distal radius fractures with a volar locking plate and screw fixation. *International Orthopedics (SCIOT)*. 2003; 27:1-6.

12. Kareem Sobky, Todd Baldini, Kenneth Thomas, Joel Bach, Allison Williams. Jennifer Moriatis Wolf, BIOMEchanical Comparison of different Volar Fracture Fixation plates for distal radius fracture. *HAND*. 2008; 3:96-101.
13. Hanae Minegishi, Osamu Dohi, Soukan An, Hidetsugu Sato. Treatment of unstable distal radius fractures with the volar locking plate. *Ups J Med Sci*. 2011; 116(4):280-284.
14. Denju Osada, Shuzo Kamei, Koichiro Masuzaki, Morimitsu Takai, Masahiro Kameda, Kazuya Tamai. Prospective study of distal radius fracture treated with volar locking plate. *Journal of Hand Surgery*. 2008, 33A.