



International Journal of Orthopaedics Sciences

E-ISSN: 2395-1958
P-ISSN: 2706-6630
IJOS 2023; 9(2): 218-221
© 2023 IJOS
<https://www.orthopaper.com>
Received: 04-03-2023
Accepted: 06-04-2023

Dr. Ismail Pandor
Junior Resident, Krishna
Institute of Medical Sciences
Karad Maharashtra

Dr. Mohit Nadkarni
Senior Resident, Krishna
Institute of Medical Sciences
Karad, Maharashtra, India

Dr. Prateek Agrawal
Junior Resident, Krishna
Institute of Medical Sciences
Karad, Maharashtra, India

Dr. Swapnil Chitnavis
Clinical Fellow, King Edward
Memorial Hospital and Seth
Gordhandas Sunderdas Medical
College, Mumbai, Maharashtra,
India

Dr. Aloukika Patil
Intern, Krishna Vishwa
Vidyapeeth, Krishna Hospital
Karad, Maharashtra, India

Corresponding Author:
Dr. Ismail Pandor
Junior Resident, Krishna
Institute of Medical Sciences
Karad, Maharashtra, India

Role of epidural analgesia in improving post-operative rehabilitation after primary definitive fixation of periarticular knee fractures followed by orthopaedic trauma

Dr. Ismail Pandor, Dr. Mohit Nadkarni, Dr. Prateek Agrawal, Dr. Swapnil Chitnavis and Dr. Aloukika Patil

DOI: <https://doi.org/10.22271/ortho.2023.v9.i2c.3373>

Abstract

Introduction: Role of epidural analgesia is well established in the literature and practice. Epidural anaesthesia is a very common procedure of anaesthesia for induction of lower limb orthopaedic cases. Considering the intra-articular fractures of the knee, they are one of the most common fractures associated with knee stiffness. Even after an adequate and rigid fracture fixation early rehabilitation is necessary to drastically improve the outcome and reduce stiffness. Hence, epidural analgesia through a lumbar epidural catheter is the most commonly used protocol in top up after surgery for post-operative analgesia. The aim of this study is to evaluate whether this method of post-operative analgesia can be helpful in improving rehabilitation after peri-articular knee injuries.

Methodology: This is a prospective study. A total of 40 patients with peri-articular of distal femur and proximal tibia were included in the study. Patients were randomised into 2 groups. Group A was given spinal anaesthesia for surgery and Group B was given combined spinal plus epidural anaesthesia with a lumbar epidural catheter. Site chosen was L1-L2 /L2-L3 for epidural analgesia and site for spinal analgesia was L3-L4 was used (in case of combined spinal epidural). Post operatively knee ROM was initiated on day 1 in both the groups. Total knee ROM was recorded on Day 2, Day 10, Day 42 (6 weeks) and Day 90 (3 months). Assessment of knee ROM was done by a single observer who was blinded for the groups.

Results: A total of 40 patients were included in this study, out of which 9 were female and 31 were males. Mean knee range of motion for group A was 20 degrees on day 2, 45 degrees at Day 10, 90 degrees at 6 weeks and 120 degrees at 3 months whereas the mean knee range of motion for group B was 40 degrees on day 2, 70 degrees at Day 10, 110 degrees at 6 weeks and 130 degrees at 3 months.

Conclusion: Peri-articular knee fractures are a common cause of limitation in knee function despite proper surgery which can be due to inadequate rehabilitation. To address this problem, it is important to give a pain-free post-operative period to the patient to develop and initiate proper post-operative rehabilitation. This can be achieved by performing peri-articular fracture fixations using Epidural anaesthesia through which post-operative analgesia can be continued in a more effective manner which also increases overall outcome of the patient.

Keywords: Epidural anaesthesia, spinal anaesthesia, post-operative rehabilitation.

Introduction

Role of epidural analgesia is well established in the literature and practice. Epidural anaesthesia is a very common procedure of anaesthesia for induction of lower limb orthopaedic cases. None the less its use post-operative after an orthopaedic procedure of lower limb is increasing in frequency. Considering the intra-articular fractures of the knee, they are one of the most common fractures associated with knee stiffness. Even after an adequate and rigid fracture fixation early rehabilitation is necessary to drastically improve the outcome and reduce stiffness. Lack of effective pain control postoperatively can significantly reduce participation in therapy regimens and thus contribute to reaccumulation of fibrotic tissue in the joint after artholysis. Hence, epidural analgesia through a lumbar epidural catheter is an established procedure to reduce post-operative pain.

1 gm preservative free morphine and 2 ml of 0.5% Bupivacaine diluted with normal saline to make a total of 10 ml is the most commonly used protocol in top up after surgery for post-operative analgesia. Epidural analgesia brings with it many honouring benefits and they include not only decrease in postoperative pain (subsequently reducing high dose opioid consumption and associated adverse effects) but also decrease in nausea and vomiting, improvement in mobilization and recovery of gastrointestinal function, decrease in length of stay (LOS), reduction in surgical stress response, and potentially, significant reduction in morbidity and mortality [1]. They are therefore commonly used to improve quality of patient care and have also been commonly used as a vital component of many enhanced recovery protocols (ERPs). It has also shown to decrease cortisol levels, expedite the return of bowel function, decrease the incidence of PE and DVT in the postoperative period, and shorten lengths of in-hospital stay [24] It may decrease the surgical risk and morbidity of certain patient populations, for example, patients with ischemic cardiac disease [25]. Despite the broad evidence base demonstrating the benefits of epidural analgesia, one must not forget to consider the adverse effects of such a technique. It also has shown to decrease post-op lung complications [25, 26]. The side effects or complications might be related to procedure or the drug used and they include dural perforation, epidural hematoma, meningitis, infection at catheter site, urinary retention, hypotension, pruritus and respiratory depression. The aim of this study is to evaluate whether this method of post-operative analgesia can be helpful in improving rehabilitation after peri-articular knee injuries.

Materials and Methods

This is a prospective study. A total of 40 patients with peri-articular of distal femur and proximal tibia were included in the study. Patients were randomised into 2 groups. Group A was given spinal anaesthesia for surgery and Group B was given combined spinal plus epidural anaesthesia with a lumbar epidural catheter. Site chosen was L1-L2 /L2-L3 for epidural analgesia and site for spinal analgesia was L3-L4 was used (in case of combined spinal epidural). The Dose for CSE (combined spinal epidural) consisted of a spinal dose of 0.5% bupivacaine heavy 3cc with fentanyl (for approximately 2-3 hr procedure) and If procedure extend beyond this time we activated the epidural with 0.5% bupivacaine 5cc + 2% lox with adrenaline 4cc diluted to make up to 10 cc. The 1st top up for epidural was given immediately post-operatively in recovery room, 2nd dose was 6-8 hours Post operatively and 3rd dose was given 12-16 hrs after the 3rd dose. Catheter was kept for a maximum of 3 days. Post operatively knee ROM was initiated on day 1 in both the groups. Total knee ROM was recorded on Day 2, Day 10, Day 42 (6 weeks) and Day 90 (3 months). Assessment of knee ROM was done by a single observer who was blinded for the groups.

Inclusion Criteria

- Distal femur and proximal tibia peri-articular fractures with previously healthy joints.
- Closed fractures or compound fractures up to Gustilo Anderson grade II.
- Adult patients of either sex.

Exclusion Criteria

- Patients with prior external fixator.
- Patients operated after 2 weeks post trauma.
- Gustilo Anderson grade III fracture.

- Previous history of knee surgery.
- History of knee arthritis, knee stiffness, affected knee ROM.
- Associated tendon, ligament, muscle injury.
- Paediatric patients
- Intra-articular unstable fractures requiring post-operative immobilization.

Results

A total of 40 patients were included in this study, out of which 9 were female and 31 were males. The mean age was 35 years. Patients were divided into 2 groups, Group A and Group B as mentioned earlier


Group A comprised of 7 distal femur fractures and 13 proximal tibia fractures. Out of 7 distal femur fractures 5 were intra-articular and 2 extra-articular fractures. 13 proximal tibia fractures were comprised of 3 extra-articular and 10 intra-articular fractures.

Group B comprised of 9 distal femur fractures and 11 proximal tibia fractures. Out of 9 distal femur fractures 6 were intra-articular and 3 were extra-articular. 11 proximal tibia fractures comprised of 2 extra-articular and 9 intra-articular fractures. Both the groups were assessed and found comparable.

Mean knee range of motion for group A was 20 degrees on day 2, 45 degrees at Day 10, 90 degrees at 6 weeks and 120 degrees at 3 months whereas the mean knee range of motion for group B was 40 degrees on day 2, 70 degrees at Day 10, 110 degrees at 6 weeks and 130 degrees at 3 months.

Table 1: Fracture, Group A and Group B

Fracture	Group A (20 Patients)	Group B (20 Patients)
Distal femur fracture		
Intra-articular	5	6
Extra-articular	2	3
Total	7	9
Proximal tibia fracture		
Intra-articular	10	9
Extra-articular	3	2
Total	13	11



Graph 1: Mean Knee Flexion (In Degrees)

Discussion

Modern anaesthesia and pre-operative medicine is commonly entangled in the challenge to create optimum treatment regimens whereby the wanted and the unwanted effects are in optimal balance in order to provide patient safety as well as comfort and to facilitate rapid recovery and rehabilitation. It is well know that surgical pain is associated with stress

responses in the intra-op period as well as post-operative period and reduction in this stress postoperatively will lead to reduced postoperative organ dysfunction and thereby provide improved outcomes [2].

Amongst the many available options for post-operative pain management, which one to choose depends on several factors such as experience of the anaesthetist, preference of anaesthetist, duration for which analgesia is required and patient preferences [3].

Very often, post-operative pain restricts the patient from initiation of rehabilitation protocol. This initial post-operative fear of pain even affects the rehabilitation after discharge because of constant fear of pain. Hence, post-operative analgesia is an important factor for boosting the confidence of the patient to gain an expected rehabilitation or range of motion post-operatively.

Data from literature shows that continuous postoperative tunnelled epidural analgesia gives promising outcomes in maintaining adequate pain control and facilitates rehabilitation postoperatively for several orthopaedic procedures including spinal surgery and adhesive capsulitis, periarticular knee fractures. Several studies in elective lower extremity orthopedic procedures have shown enhanced rehabilitation by post-operative regional analgesia techniques. Continuous epidural analgesia has also shown to be a viable option for managing persistent stiffness after TKA [4]. Continuous infusion of local anaesthetic or opioid alone or in combination provide effective postoperative pain relief but the most effective ones alone or in combination, dose and route remain controversial [5]. Wu *et al.* compared the effectiveness of systemic opioids, epidural opioid and epidural opioid-local anesthetic mixture. They found that epidural opioid was more effective than systemic opioids [6].

In a meta-analysis on epidural versus systemic analgesia in various surgical procedures, epidural local anesthetic were found to reduce postoperative complications compared with epidural or systemic opioid techniques [7]. Postoperative epidural analgesia has been shown to decrease cardiac ischemia post operatively and postoperative epidural analgesia has been shown in a meta-analysis of a variety of surgical procedures to reduce the incidence of postoperative pulmonary complications. Local epidural anaesthesia have also shown reduction in post-operative paralytic ileus and improved post-operative cognitive function in elderly patients in various surgical procedures [8-11].

Despite its widespread benefits epidural analgesia carries potential risks such as dural perforation, epidural hematoma and infection etc. [20] Chaney MA [21] reported classical side effects of epidural opioids being pruritis, nausea, vomiting, urinary retention and respiratory depression. Nonetheless he found that majority of them were dose dependant. A study by Stenseth *et al.* [22] using epidural morphine found that 90 percent were completely satisfied with the course. Despite the positive results side effects like pruritis (11%), nausea and vomiting (34%) and respiratory depression (0.9%) was seen. Similarly Shafiq *et al.* [23] in one study found that over all incidence of complications was 26.6 percent. Common complications were motor block, dural tap, accidental catheter pullouts, ineffective pain control etc. Nonetheless our study did not show any such complications.

There are certain limitations to the study. This study included combined patients of peri-articular knee fractures with both intra-articular as well as extra-articular. A more selective study is required for strengthening the study.

Conclusion

Peri-articular knee fractures are a common cause of limitation in knee function despite proper surgery which can be due to inadequate rehabilitation. To address this problem, it is important to give a pain-free post-operative period to the patient to develop and initiate proper post-operative rehabilitation. This can be achieved by performing peri-articular fracture fixations using Epidural anaesthesia through which post-operative analgesia can be continued in a more effective manner which also increases overall outcome of the patient.

Conflict of Interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Financial Support

Not available

References

1. Choi S, Trang A, McCartney C. Reporting functional outcome after knee arthroplasty and regional anesthesia: A methodological primer. *Reg Anesth Pain Med* 2013;38:340-9.
2. Kehter IL. Multimodal approach to control postoperative I Pathophysiology and rehabilitation. *Br. J Anaesth.* 1997;78:606-17
3. Savaye C, McQuitty C, Wang D *et al.* Post Thoracotomy pain management. *Chest Surg Clin Nam,* 2002;12(2):251-63.
4. Pariente GM, Lombardi AV Jr, Berend KR, Mallory TH, Adams JB. Manipulation with prolonged epidural analgesia for treatment of TKA complicated by arthrofibrosis. *Surg Technol Int.* 2006;15:221-224.
5. Glass PSA, Stock P, Ginsberg B, Goldberg JS, Slide ON. Use of patient controlled analgesia to compare the efficacy of epidural to intravenous fentanyl administration: *Anesth Analg.* 1992;74:345-351.
6. Wu CL, Cohen SR, Richman JM, *et al.* Efficacy of postoperative patient-controlled and continuous infusion epidural analgesia versus intravenous patient-controlled analgesia with opioids. *Meta-analysis. Anesth.* 2005;103(5):1079-83.
7. Ballantyne JC, Carr DB, de Ferranti S, Suarez T, Lau J, Chalmers TC, Angelillo I, Mosteller F. The comparative effects of postoperative analgesic therapies on pulmonary outcome. Cumulative meta-analysis of randomized controlled trials. *Anesth Analg.* 1998;86:598-612.
8. Kehlet H, Holte K. Effect of postoperative analgesia on surgical outcome, *Br J Anaesth.* 2001;87:62-72.
9. Veering BT, Cousins MJ - Cardiovascular and pulmonary effects of epidural anaesthesia. *Anaesth Intens Care.* 2000;28:620-635.
10. Curli E, Trudel J, Belliveau P. Return of bowel function after thoracic epidural anesthesia. *Dis Colon Rectum;* c2001.
11. Hijorto NC, Neumann P, Frosig F, *et al.* A controlled study on the effect of Epidural analgesia with local anaesthetics and morphine on morbidity after abdominal surgery. *Acta Anaesthesiol Scand.* 1985;29:790-6.
12. Jayr C, Thomas H, Rey A, Farhat F, Lasser P, Bourgain JL. Postoperative pulmonary complications, Epidural analgesia using bupivacaine and opioid versus parenteral opioid. *Anesthesiology.* 1993;78:666-76.
13. Liu S, Carpenter RD, Mackey DC, *et al.* Effects of

- postoperative analgesic technique on rate of recovery after colon surgery *Anesthesiology*. 1995;83:757-65.
14. Mann C, Pouzeruttee Y, Boccara G, *et al.* Comparison of intravenous or epidural patient-controlled analgesia in the elderly after major abdominal surgery *Anesthesiology* 2000;92:433-41.
 15. Seeling W, Bothner U, Eifert B, *et al.* Patient-controlled analgesia versus epidural analgesia using bupivacaine and morphine following major abdominal surgery. No difference in postoperative morbidity *Anaesthesist*. 1991;40:614-23.
 16. Seeling W, Bruckmooser KP, Hufner C, Kneitingner E, Ragu C, Ruckemar M. No reduction in postoperative complications by the use of catheterized epidural analgesia following major abdominal surgery. *Anaesthesist*. 1990;39:33-40.
 17. Holte K, Kehlet H. Epidural anesthesia and risk of anastomotic leakage. *Reg Anesth Pain Med*. 2001;26:111-7.
 18. Christopherson R, Beattie C, Frank SM, *et al.* Perioperative Ischemic Preconditioning Study Group Perioperative morbidity in patients randomized to epidural or general anesthesia for lower extremity vascular surgery. *Anesthesiology*. 1993;79:422-34.
 19. Yeuger MP, Glass DD, Nell RK, Brinck-Johnsen T. Epidural anesthesia and analgesia in high risk surgical patients. *Anesthesiology*. 1987;66:729-36.
 20. Andreas M, Zollinger A, De Lorenzi D, *et al.* Prospective, randomized comparison of extrapleural versus epidural analgesia for post thoracotomy pain. *Ann Thorac Surg*. 1998 66:367-72.
 21. Chaney MA. Side effects of intrathecal and epidural opioids. *Can J Anaesth*. 1995 Oct;42(10):891-903.
 22. Stenseth R, Sellevold O, Breivik H. Epidural morphine for postoperative pain: Experience with 1085 patients. *Acta Anaesthesiol Scand*. 1985 Jan;29(1):148-56.
 23. Shafiq FI, Hamid M, Samud K. Complications and interventions associated with epidural analgesia for postoperative pain relief in a tertiary care hospital. *Middle East J Anaesthesiol*. 2010 Oct;20(6):827-32.
 24. Strandness T, Wiktor M, Varadarajan J, Weisman S. Migration of pediatric epidural catheters. *Paediatr Anaesth*. 2015 Jun;25(6):610-3.
 25. Gerheuser F, Roth A. [Epidural anesthesia]. *Anaesthesist*. 2007 May;56(5):499-523; quiz 524-6.
 26. Moriarty A. Pediatric epidural analgesia (PEA). *Paediatr Anaesth*. 2012 Jan;22(1):51-5.

How to Cite This Article

Pandor I, Nadkarni M, Agrawal P, Chitnavis S, Patil A. Role of epidural analgesia in improving post-operative rehabilitation after primary definitive fixation of periarticular knee fractures followed by orthopaedic trauma. *International Journal of Orthopaedics Sciences*. 2023;9(2):218-221.

Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.